

Osteochondral Autograft Transfer (OATs)

Postoperative Instructions

MEDICATIONS: You will be given three prescriptions- Medications to control pain, nausea, and constipation. Take as directed. You should also eat yogurt for the first 3 days to help maintain the normal, helpful bacteria within your stomach.

DRESSING: Leave your dressing on until 48 hours after surgery. At 48 hours after surgery, you may remove your dressing. **LEAVE THE TAPE STRIPS OVER YOUR INCISIONS.** These will stay on for 1 ½- 2 weeks and will slowly peel off.

SHOWERING: You may shower 48 hours after your surgery and dry your incisions. Do **NOT** immerse in a tub or pool for 4 weeks to avoid excessive scarring and risk of infection.

ICE: Icing is very important to decrease swelling and pain and improve mobility. During the first 24 hours, apply ice every 2 hours, 15 minutes each time. Or, if you received an ice machine, use at all times over a towel. After 24 hours, continue to use the ice 3 – 4 times a day, 15 – 20 minutes each time to keep swelling to a minimum.

WEIGHT BEARING: You should not bear weight on your leg until cleared by your surgeon. Use your crutches for ambulation.

ACTIVITIES: Rest and elevate your leg for the first 24 hours. Do **NOT** place a pillow under your knee. Elevate with a pillow under your calf and ankle. It is important to get your leg as straight as possible as soon as possible.

EXERCISES:

1) Quadri-cep Contractions:	5 – 10 per hour Push knee into bed Tighten thigh muscle
2) Straight Leg Raises:	5 per hour
3) Ankle pumps	10 per hour

If you experience severe pain that your pain medication does not relieve, you should let us know. **If you experience a temperature over 101.5*, redness or swelling in your thigh or calf, please contact our office daytime, evenings or weekends at (847) 634-1766.**

GOALS: Safe ambulation and pain control.