



# Proximal Hamstring Repair

**MEDICATIONS:** You will be given three prescriptions- Medications to control pain, nausea, and constipation. Take as directed. You should also eat yogurt for the first 3 days to help maintain the normal, helpful bacteria within your stomach.

**DRESSING:** You may remove your dressing 48 hours after surgery. You do not need to reapply a dressing unless you would like. **LEAVE THE TAPE STRIPS OVER YOUR INCISIONS.** These will stay on for 1 ½- 2 weeks and will slowly peel off.

**SHOWERING:** You may shower after your dressing is changed letting soap and water run over the incision, no scrubbing. You do not need to redress the incisions. Do **NOT** immerse in a tub or pool for 4 weeks to avoid excessive scarring and risk of infection.

**ICE:** Icing is very important to decrease swelling and pain and improve mobility. During the first 24 hours, apply ice every 2 hours, 15 minutes each time. After 24 hours, continue to use the ice 3 – 4 times a day, 15 – 20 minutes each time to keep swelling to a minimum.

**WEIGHT BEARING:** You should not put weight on your leg until cleared by your doctor. Use your crutches until your first therapy visit. Maintain the hinged knee brace per the protocol. Contact your doctor if you would like use a knee scooter.

**ACTIVITIES:** Rest your leg for the first 24 hours. Do not elevate it unless specifically allowed.

**EXERCISES:**      1) Ankle pumps                      10 per hour

If you experience severe pain that your pain medication does not relieve, you should let us know. **If you experience a temperature over 101.5°, redness or swelling in your thigh or calf, please contact our office daytime, evenings or weekends at (847) 634-1766.**

**GOALS:** Pain control within 1-2 weeks